

Assessment of Anxiety Related To Osce Exam Among Alzaeim Al-Azahari University Medical Students 2022

Saja Abd Alrhaman Alshreef¹, Mohammed Hammad Jaber^{1,*}, Solafa Othman Elbadawi¹,
Muhammad Ashir Shafique², Sara Salah Eldin Ali¹, Hiba Salah Abdelgadir¹

¹Faculty of Medicine, Al-Zaeim Al-Azhari University, Khartoum, Sudan

²Department of Medicine, Jinnah Sindh Medical University, Karachi, Pakistan

Email address:

mohammesjaber123@gmail.com (Mohammed Hammad Jaber)

*Corresponding author

To cite this article:

Saja Abd Alrhaman Alshreef, Mohammed Hammad Jaber, Solafa Othman Elbadawi, Muhammad Ashir Shafique, Sara Salah Eldin Ali et al. (2023). Assessment of Anxiety Related To Osce Exam Among Alzaeim Al-Azahari University Medical Students 2022. *American Journal of Health Research*, 11(6), 158-163. <https://doi.org/10.11648/j.ajhr.20231106.12>

Received: November 1, 2023; **Accepted:** November 16, 2023; **Published:** December 18, 2023

Abstract: Introduction: This study to gather information about the Incidence rate of severe anxiety before and during OSCE exam among medical students and detect the risk factors for it, in order to increase the awareness level of medical students' community about the effects of anxiety before and during exams on their academic achievement. Methodology: a cross-sectional study was done in medical students at AAU in fifth year The results: A total 211 sample was collected, 48.3% were male while 51.7% were female. 53.6% of participants age from 20 to 23 while 46.4% age from 24 to 26. 79.6% of participants believe that the anxiety and fear caused by your fear of forgetting the information, while 20.4% of participants don't believe that. 75.4% of participants believe that anxiety because of the lot to comprehend and less time to practice while 24.6% don't believe that. 93.8% of participants get an anxiety episode because of the examiner and how he is treating participants, Conclusion: 97.2% of students reported feeling anxious throughout the OSCE exam, There was no collaboration between the examiner and the student which made students more anxious. The time in not enough for every station. We recommended students to adopt healthy & effective mean of relaxation like yoga, breathing exercises and non formal counseling.

Keywords: Anxiety, Osce Exam, Medical Student, Risk Factors

1. Introduction

Anxiety disorders are real, serious medical conditions - just as real and serious as physical disorders such as heart disease or diabetes. Anxiety disorders are the most common and pervasive mental disorders in the United States [1].

Generalized Anxiety Disorder (GAD) is characterized by persistent and excessive worry about several different things. People with GAD may anticipate disaster and may be overly concerned about money, health, family, work, or other issues. Individuals with GAD find it difficult to control their worry. They may worry more than seems warranted about actual events or may expect the worst even when there is no apparent reason for concern.

GAD is diagnosed when a person finds it difficult to control worry on more days than not for at least six months and has

three or more symptoms. This differentiates GAD from worry that may be specific to a set stressor or for a more limited period.

GAD affects 6.8 million adults, or 3.1% of the U.S. population, in any given year. Women are twice as likely to be affected. The disorder comes on gradually and can begin across the life cycle, though the risk is highest between childhood and middle age. Although the exact cause of GAD is unknown, there is evidence that biological factors, family background, and life experiences, particularly stressful ones, play a role.

Sometimes just the thought of getting through the day produces anxiety. People with GAD don't know how to stop the worry cycle and feel it is beyond their control, even though they usually realize that their anxiety is more intense than the situation warrants. All anxiety disorders may relate to a difficulty tolerating uncertainty and therefore many people with GAD try to plan or control situations. Many people

believe worry prevents bad things from happening, so they view it is risky to give up worry. At times, people can struggle with physical symptoms such as stomachaches and headaches.

When their anxiety level is mild to moderate or with treatment, people with GAD can function socially, have full and meaningful lives, and be gainfully employed. Many with GAD may avoid situations because they have the disorder or they may not take advantage of opportunities due to their worry (social situations, travel, promotions, etc.). Some people can have difficulty carrying out the simplest daily activities when their anxiety is severe.

Test anxiety has been defined as the emotional, physiological, and behavioral responses surrounding the potential consequences of negative evaluation on an upcoming test or exam. Recent work by Lowe and colleagues suggests that biopsychosocial factors, including biological (e.g., physiological arousal), psychological (e.g., emotional or cognitive factors), and social (e.g., parent pressure), contribute to development and expression of test anxiety. In association with these factors, students with test anxiety exhibit interfering thoughts, feelings, or off-task behaviors in testing situations that are perceived to be personally relevant.

Additionally, when a test-anxious student perceives a test to be threatening and performs less than optimally, that student's maladaptive cognitions are reinforced and test anxiety increases. It has been estimated that between 10% and 40% of all students suffer from various levels of test anxiety, and students with disabilities, women, and minority students report higher rates of test anxiety. Test anxiety has been reported to onset as early as age 7, and when present, test anxiety is a disruptive factor in students' academic careers. Students with high levels of test anxiety perform lower on tests and have lower overall academic achievement as measured by grade point averages [2].

Medical schools around the world aim to train and produce competent and empathetic physicians to help the sick, advance medical knowledge, and promote public health. However, medical education is one of the most academically and emotionally demanding training programs out of any profession, and consequently, the time and emotional commitment necessary for medical students to devote to their training is extensive. Such demands and stress cause a negative effect on the students' psychological well-being, and can precipitate is often undetected and undertreated in the general population. In addition to intense feelings of fear or panic, sufferers of anxiety can experience other physiological symptoms including fatigue, dizziness, headaches, nausea, abdominal pain, palpitations, shortness of breath, and urinary incontinence.

Anxiety can also impair goal-directed attention and concentration, working memory, and perceptual-motor function, all of which are important domains which enable medical students and physicians to provide safe and efficacious medical care to patients [3].

The OSCE was a most anxiety-provoking assessment method and students prepared more for the OSCE than for the other examinations. The expectation to succeed was also higher for the OSCE [6-9]. State anxiety during the OSCE

was associated with the level of preparation but not with scores obtained. Medical education exposes students to a highly stressful environment. Numerous reasons have been attributed, however OSCE and examinations are amongst the top stressors related to anxiety being experienced by the graduating students. Here on Sudan, there is no reliable data and reports about the association of test anxiety on OSCE performance. So This study is necessary to gather information about the Incidence rate of severe anxiety before and during OSCE exam among medical students and detect the risk factors for it, in order to increase the awareness level of medical students' community about the effects of anxiety before and during exams on their academic achievement.

2. Methodology

2.1. Study Design

This study followed descriptive cross sectional study.

2.2. Study Area

The study was conducted in Al- Zaem Al -Azhari University. Bahri Locality, Khartoum state, Sudan.

2.3. Study Duration

This study was conducted from December 2021- June 2022.

2.4. Study Population

All Medical students at AAU in fifth year which all about 211.

2.5. Inclusion Criteria

1. All Medical students at AAU in fifth year.
2. Those who agreed to participate in the study.

2.6. Exclusion Criteria

Those who refused to participate in the study.

2.7. Sampling Technique

Simple random sampling method was used to select the sample size.

2.8. Sample Size

The sample size was calculated by the following formula at the confidence level of 95% and degree of precision 0.05.

$$n = \frac{N}{1+N(d^2)}$$

$$n=212/[1+(0.000125)]$$

n= sample size.

N = Population Size = Medical college student

d = degree of precision

so, n=211.97

2.9. Data Collection Tool and Technique

Closed questionnaire was designed to provide information, the collection was done by sending the questionnaire in special whatsapp groups.

2.10. Study Variables

Sex || distribution: males and females.

Age || of the student

Personal || habits: smoking, alcohol, drugs, or coffee.

Average || reading hours.

History of mental illness or family history of mental illness. Current medications.

2.11. Data Analysis and Interpretation

The data was reviewed, ordered, and entered manually into a computer and then analyzed using Statistical Package for the Social Sciences (SPSS) version 25.0. Data was presented as frequencies (%) and mean values (SD) as appropriate and compared using either the chi-square test (for proportions) or the student's T-test for mean values. $P < 0.05$ as statistically significant.

3. Ethical Issues


1. Ethical approval was obtained from Ethics and Research Committees in the AAU.
2. Permission from the students was taken.
3. Students were informed that their data privacy will be considered and used for the purpose of the study only.

4. The Results

A total 211 sample was collected, 48.3% were male while 51.7% were female. 53.6% of participants age from 20 to 23 while 46.4% age from 24 to 26.

Table 1. Distribution of participants according to their age and gender ($n=211$).

| Variables | Frequency | Percent |
|---------------|-----------|---------|
| Gender | | |
| Male | 102 | 48.3 |
| Female | 109 | 51.7 |
| Age | | |
| From 20 to 23 | 113 | 53.6 |
| From 24 to 26 | 98 | 46.4 |



97.2% of participants feel anxious during OSCE while 2.8% don't feel that

Figure 1. Distribution of participants according to their feeling of anxiety ($n=211$).




Figure 2. Distribution of participants according to their feeling more anxious than other colleagues before the exam ($n=211$).

53% of participants don't feel anxious more than other colleagues while 47% of participants do feel that.

Table 2. Distribution of participants according to their causes to anxiety ($n=211$).

| Variables | Frequency | Percent |
|--|-----------|---------|
| Is the anxiety and fear caused by your fear of forgetting the information? | | |
| Yes | 168 | 79.6 |
| No | 43 | 20.4 |
| Is the anxiety because of the lot to comprehend and less time to practice? | | |
| Yes | 159 | 75.4 |
| No | 52 | 24.6 |

79.6% of participants believe that the anxiety and fear caused by your fear of forgetting the information, while 20.4% of participants don't believe that. 75.4% of participants believe that anxiety because of the lot to comprehend and less time to practice while 24.6% don't believe that.


Table 3. Distrubtion of participants according to the effect of anxiety (n=211).

| Variables | Frequency | Percent |
|--|-----------|---------|
| Do you get an anxiety episode because of the examiner and how will he treat you? | | |
| Yes | 198 | 93.8 |
| No | 13 | 6.2 |
| Do you feel headache before or during the exam? | | |
| Yes | 47 | 22.3 |
| No | 164 | 77.7 |
| Do you feel anxious just thinking you don't have time? | | |
| Yes | 151 | 71.6 |
| No | 60 | 28.4 |
| Do you sweat a lot from your hands and face during the exam? | | |
| Yes | 75 | 35.5 |
| No | 136 | 64.5 |
| Do you get a gastrointestinal symptoms before the exam? | | |
| Yes | 122 | 57.8 |
| No | 89 | 42.2 |


93.8% of participants get an anxiety episode because of the examiner and how he treating participants, while 6.2% don't get it. 22.3% of participants feel headache before or during the exam while 77.7% don't feel. 71.6% of participants feel anxious just thinking you don't have time, while 28.4% don't feel that. 35.5% of participants do sweat a lot from your hands and face during the exam, while 64.5% don't. 57.8% of participants get a gastrointestinal symptoms before the exam, while 42.2% don't get.

Table 4. Distrubtion of participants according to the effect of anxiety on sleeping (n=211).


| Variables | Frequency | Percent |
|---|-----------|---------|
| Does anxiety affect your sleep? | | |
| Yes | 144 | 68.2 |
| No | 67 | 31.8 |
| Have you ever felt the need to use sleeping pills to sleep? | | |
| Yes | 23 | 10.9 |
| No | 188 | 89.1 |

**Figure 3.** Distrubtion of participants according to effect of anxiety on exam performance (n=211).

67% of participant's academic performance have affected by anxiety while 33% have not.

**Figure 4.** Anxiety getting relieved with the recurrence of exams (n=211).

60% of participant's anxiety getting relieved with the recurrence of exams while 40% are not.



64% of participants satisfy with results while 36% don't.

Figure 5. Distrubtion of participants according to there satisfaction of the results (n=211).

5. Discussion

In our study males and females were nearly equal (51.7%), (48.3%) the same result was also report in Korea by Kyong-Jee Kim et al [4] and other's studies by B. Pahwa et al. and Majumder MA et al demonstrated that thier is more females than males [2, 4] but males seem to have been slightly more

prevalent than females, according to Zhang N. et al [10].

According to the previous studies, the mean age of the students was 24.4 [10] and 24.2 (5) 20.1 (21) and this nearly same as our results that the age group 20 _23 comprised the maximum number, but a higher mean age 25.9 was reported in another study. [3] and less mean also reported in previous study [14].

Niu Zhang noted that 85% of students reported feeling

anxious throughout the OSCE exam, which is lower than our study's 97.2%. [10] Khalid I. Khoshhal et al found a prevalence of exam anxiety in Saudi Arabia of 65%, which is also lower than our findings. [12] In Pakistan Brand and Schoonheim - Klein et al reported greater stress levels among dental students during OSCE in comparison to written and practical assessment methods, they said students had to prepare better for OSCEs and their expectation to pass OSCE was also significantly higher which might be the cause of stress [18] In another study students said that there is a need to reassess the amount of the study materials allocated in their study and this could be the cause of their anxiety [12].

A variety of factors can contribute to assessment stress and anxiety among medical students, physiological factors are considered a leading cause of assessment anxiety [5], Mavis found that students with more clinical experience about the clinical exams, and he suggested that as students gain more clinical experience they become more knowledgeable about what they know and what they don't know, and the latter can increase anxiety relative to students with less clinical experience. [3] Niam H, et al also observed several factors that contribute to assessment anxiety among final year medical students, extensive course load, the long duration of periods of assessment, lack of assessment preparation and lack of exercise were reported to be the most important contributing factors [5].

In the current study 79.6% of the students believed that anxious is main reason for their forgetting and this affects their performance, [30] in Taif Medical College they conducted the anxiety has a negative effect on overall academic performance of Medical students, and they reported that stress was more apparent in the third year. [17], anxiety was not predictive of performance outcome in the OSCE, written examination or preclinical preparation test [18].

In a previous study 42% reported that this stress did not affect their performance while decrease in performance was reported by 33% (21) another study reported the same result [1] but there's another study reported that anxiety and self-efficacy were negatively associated: students who were highly anxious about the OSCE were less confident about their ability to perform in the stimulation [3].

Two third of the participants because of the lot comprehend and less time and this also report by other studies when asked students about their comments they said that they need more time and Majumder MD et al reported the same result [2].

Regarding the examiners and how they interacted with the students, we noticed that they were cooperative, that there was a good staff available, and that there was a good mixture of local and foreign examiners [2]. However, because of the examiner's, 93.8% who study become anxious [11-16].

Many students worry that they don't have enough time, and respondents also thought that the time allotted for projects was insufficient, which added to their stress [19-22]. However, there is disagreement among students and examiners in the same study regarding the timing at OSCE stations [2].

A prior survey revealed that the majority of students were satisfied with their OSCE performance as indicated in 64 percent of the participants said they were satisfied with their results [23-26], many students also had little knowledge of assessment taking and anxiety reduction techniques [27-29].

6. Conclusion

In our study, we assessed the anxiety related to OSCE Exam in Medical Students of AAU we found that

1. There was a very high prevalence of anxiety in OSCE exams among students (97%)
2. Equal percent between males and females
3. Anxiety decreased students performance because of their stress.
4. There was no collaboration between the examiner and the student which made students more anxious.
5. The time in not enough for every station.

7. Recommendation

Due to the significant frequency of mental distress among students,

1. we advise policy-makers to pay attention to the issue and take appropriate corrective measures. representatives from the college, non-governmental organizations, parents, students, and other relevant individuals. These recognized causes of mental distress against students must be addressed in programs designed to prevent it.
2. Specialized intervention be offered to those with high anxiety levels.
3. Encourage students to adopt healthy & effective mean of relaxation like yoga, breathing exercises and non formal counseling.

ORCID

Mohammed Hammad Jaber: <https://orcid.org/0000-0002-1366-2375>

Conflicts of Interest

The authors declare no conflicts of interest.

References

- [1] Test anxiety. Anxiety and Depression Association of America. www.adaa.org/living-with-anxiety/children/test-anxiety. Accessed July 28, 2014.
- [2] Von der Embse, Nathaniel & Barterian, Justin & Segool, Natasha. (2013). Test Anxiety Interventions for Children and Adolescents: A Systematic Review of Treatment Studies from 2000–2010. *Psychology in the Schools*. 50. 57-71. 10.1002/pits.21660.

- [3] Quek TT, Tam WW, Tran BX, Zhang M, Zhang Z, Ho CS, Ho RC. The Global Prevalence of Anxiety Among Medical Students: A Meta-Analysis. *Int J Environ Res Public Health*. 2019 Jul 31; 16(15): 2735.
- [4] Jay A. Students' perceptions of the OSCE: a valid assessment tool?. *British Journal of Midwifery*. 2007 Jan; 15(1): 32-7.
- [5] Majumder MA, Kumar A, Krishnamurthy K, Ojeh N, Adams OP, Sa B. An evaluative study of objective structured clinical examination (OSCE): students and examiners perspectives. *Advances in medical education and practice*. 2019; 10: 387.
- [6] Mavis B. Self-efficacy and OSCE performance among second year medical students. *Advances in health sciences education*. 2001 May; 6(2): 93-102.
- [7] Kim KJ. Factors associated with medical student test anxiety in objective structured clinical examinations: a preliminary study. *International journal of medical education*. 2016; 7: 424.
- [8] Hashmat S, Hashmat M, Amanullah F, Aziz S. Factors causing exam anxiety in medical students. *Journal-Pakistan Medical Association*. 2008 Apr 1; 58(4): 167.
- [9] Pendergast L, Watkins M, Canivez G. Educational Psychology: An International Journal of Experimental Educational Psychology.
- [10] Lyndon MP, Strom JM, Alyami HM, Yu TC, Wilson NC, Singh PP, Lemanu DP, Yelder J, Hill AG. The relationship between academic assessment and psychological distress among medical students: a systematic review. *Perspectives on medical education*. 2014 Dec; 3(6): 405-18.
- [11] Vasli P, Shahsavari A, Estebarsari F, AsadiParvar-Masouleh H. The predictors of nursing students' clinical competency in pre-internship objective structured clinical examination: The roles of exam anxiety and academic success. *Nurse Education Today*. 2021 Dec 1; 107: 105148.
- [12] Andika AA, Susani YP, Sari DP. FACTORS CORRELATED WITH MEDICAL STUDENTS ANXIETY IN CLINICAL SKILLS EXAMINATION. *Jurnal Pendidikan Kedokteran Indonesia: The Indonesian Journal of Medical Education*; 10(3): 215-26.
- [13] Zhang N, Henderson CN. Test anxiety and academic performance in chiropractic students. *Journal of Chiropractic Education*. 2014; 28(1): 2-8.
- [14] Pahwa B, Goyal S, Srivastava K, Saldanha D, Bhattacharya D. A study of exam related anxiety amongst medical students. *Industrial Psychiatry Journal*. 2008 Jan 1; 17(1): 46.
- [15] Khoshhal KI, Khairy GA, Guraya SY, Guraya SS. Exam anxiety in the undergraduate medical students of Taibah University. *Medical teacher*. 2017 Mar 16; 39(sup1): S22-6.
- [16] Guraya SY, Guraya SS, Habib F, AlQuiliti KW, Khoshhal KI. Medical students' perception of test anxiety triggered by different assessment modalities. *Medical teacher*. 2018 Jul 6; 40(sup1): S49-55.
- [17] Colbert-Getz JM, Fleishman C, Jung J, Shilkofski N. How do gender and anxiety affect students' self-assessment and actual performance on a high-stakes clinical skills examination?. *Academic medicine*. 2013 Jan 1; 88(1): 44-8.
- [18] Rupani MP, Parikh KD, Trivedi AV, Singh MP, Shah K, Parmar K, Shakya K, Chauhan K, Ganveet J. Cross-sectional study on exam anxiety among medical students of a tertiary care teaching hospital of western India. *Natl J Community Med*. 2016; 7(5): 449-54.
- [19] Naim H, Khan WA, Naim HH, Jawaaid M. Examination related anxiety and its management amongst medical students. *Journal for Studies in Management and Planning*. 2014; 1: 104-14.
- [20] Arain FR. Does test anxiety and gender have an impact on OSCE Performance among medical students?. *Middle East Journal of Family Medicine*. 2021 Nov 1; 7(10): 101.
- [21] Brand HS, Schoonheim-Klein M. Is the OSCE more stressful? Examination anxiety and its consequences in different assessment methods in dental education. *European Journal of Dental Education*. 2009 Aug; 13(3): 147-53.
- [22] Naim H, Khan WA, Naim MH, Jawaaid M. Exam related anxiety and its management amongst medical students.
- [23] Hadi MA, Ali M, Haseeb A, Mohamed MM, Elrggal ME, Cheema E. Impact of test anxiety on pharmacy students' performance in Objective Structured Clinical Examination: a cross-sectional survey. *International Journal of Pharmacy Practice*. 2018 Apr; 26(2): 191-4.
- [24] Mittal R, Kumar R. Exam stress in MBBS students and the methods used for its alleviation. *International Journal of Medical and Dental Sciences*. 2018 Jan 9: 1604-8.
- [25] Kurt AS, Balci S, Kose D. Test anxiety levels and related factors: Students preparing for university exams. *J Pak Med Assoc*. 2014 Nov 1; 64(11): 1235-9.
- [26] Bedaso A, Duko B, Yeneabat T. Predictors of mental distress among undergraduate health science students of Hawassa University, College of Medicine and Health Sciences, Hawassa, SNNPR, Ethiopia: a cross-sectional study. *Annals of general psychiatry*. 2020 Dec; 19(1): 1-5.
- [27] Ferreira ÉD, Pinto RZ, Arantes PM, Vieira ÉL, Teixeira AL, Ferreira FR, Vaz DV. Stress, anxiety, self-efficacy, and the meanings that physical therapy students attribute to their experience with an objective structured clinical examination. *BMC Medical Education*. 2020 Dec; 20(1): 1-9.
- [28] Kavakci O, Semiz M, Kartal A, Dikici A, Kugu N. Test anxiety prevalence and related variables in the students who are going to take the university entrance examination. *Dusunen Adam The Journal of Psychiatry and Neurological Sciences*. 2014; 27(4): 301.
- [29] Inam SN, Saqib A, Alam E. Prevalence of anxiety and depression among medical students of private university. *Journal-Pakistan Medical Association*. 2003 Feb 1; 53(2): 44-6.
- [30] Von Der Embse N, Barterian J, Segool N. Test anxiety interventions for children and adolescents: A systematic review of treatment studies from 2000–2010. *Psychology in the Schools*. 2013 Jan; 50(1): 57-71.